


*Liquid
Bio-Fertilizer
for Roots*



SuperRoots

With SuperTrace, Nutrex-M/F, SoilSeeds, Mycorrhizae & Humic Substances*

For distribution in the State of Oregon

5-0-6 – 2% Naturally Chelated Iron

No Phosphates - No EDTA

*The humic substances are derived from ancient shale, the most thoroughly composted vegetation on Earth.

SuperRoots™ is a proprietary suspended concentrate bio-fertilizer for tree and shrub root treatments and turf application after aeration. **SuperRoots™** contains **Nutrex-M/F™** stabilized macro-nutrients, **SuperTrace™** fully chelated micro-nutrients, endo-ecto mycorrhizae with trichoderma/scleroderma root shield fungi, and **SoilSeeds™** plant-derived microbes and humic substances (USDA compliant). **SuperRoots™** is not toxic to plants when used as directed. Biological activity may occur (float on the surface) when stored for extended times. If this occurs, in-line screens should be used to prevent nozzle clogging.

To protect from degradation, do not store in sunlight or near heat source.

Net contents: 2.5 gal Net weight: 23.63 lb Density: 9.45 lb/gal Batch Code: 071213SR Expiration: 07-12-16



3944 South Bay Rd NE, Olympia, WA 98516 • 360.491.0556 • www.plantessentials.biz

A Portion of the Proceeds from this Product Goes to Support the Northwest Parkinson's Foundation – www.nwpcf.org

Guaranteed Analysis (By Weight)

Nitrogen (N)	5.00%
5.00% Urea Nitrogen	
Soluble Potash (K ₂ O)	6.00%
Iron (Fe)	2.00%
2.00% Chelated Iron	
Manganese (Mn)	0.125%
0.125% Chelated Manganese	
Molybdenum (Mo)	0.001%
Zinc (Zn)	0.050%
0.050% Chelated Zinc	

Derived from: Urea, potassium hydroxide, sodium molybdate, chelated ferrous sulfate, chelated manganese sulfate, and chelated zinc sulfate.

Contains non-plant food ingredients: Endo/ecto mycorrhizae with trichoderma/scleroderma root shield fungi: Glomus intraradices (625 propagules), G. mosseae (625 propagules), G. aggregatum (625 propagules), G. etunicatum (625 propagules), Rhizopogon villosulus (1.25 million propagules), R. luteolus (1.25 million propagules), R. amylopogon (1.25 million propagules), R. fulvigleba (1.25 million propagules), Pisolithus tinctorius (2.5 million propagules), Scleroderma cepa (1.25 million propagules), S. citrinum (1.25 million propagules) Trichoderma konigii (1.8 million cfu) and T. harzianum (1.8 million cfu) Dicyandiamide 0.16% N-(n-butyl) thiophosphoric triamide ammoniacal nitrogen additives 0.0084% Lactobacillus perans 11,800 CFU/ml Lactobacillus paracasei 11,800 CFU/ml Ascophyllum nodosum from brown seaweed 0.0046% Humic Substances (derived from Leonardite) 0.013 %*
*analyzed by CDFA method

Information regarding the contents and levels of metals in this product is available on the internet at: <http://www.aapfco.org/metals.htm>

Tank mix with water and apply to soil. No other additives needed.

FIELD GROWN NURSERY STOCK:

Apply at the rate of 2-5 gal of **SuperRoots™** per acre in 20-200 gal of water. Immediately irrigate following application to soil. **DO NOT APPLY TO FOLIAGE.** In case of accidental foliage contact, rinse with water immediately.

TREES & SHRUB ROOT FERTILIZING:

Surface application: Mix 2.5 gal of **SuperRoots™** with 10 gal of water per 1,250 sq ft. Apply with a shower-head gun to the soil over plant roots. **DO NOT APPLY TO FOLIAGE.** In case of accidental foliage contact, rinse with water immediately. For best results, apply in spring and fall. Mulch will inhibit movement of this product to the soil. Apply directly to soil for best results.

Sub-soil injection (moist soil only): Mix 2.5 gal of

SuperRoots™ with 10 gal of water. Inject 2 oz of diluted **SuperRoots™** per injection site (based on using 200# pressure with a 3/8" hose), 1-4" deep in a 15 inch grid pattern from the stem or trunk to 2.5 ft beyond the drip line. Start injections under any mulch. Do not apply more than 4" deep in soil. **Dry soil should be watered to a depth of 6" before treatment.**

Treating roots under grass: Treating roots in turf requires deeper injection because of root competition and bacterial/fungal antagonism. Inject **SuperRoots™** below the grass roots where feasible. Ideally, grass cover should be removed and certified pathogen free compost installed to provide the best conditions for healthy plant root development.

WATERING:

Watering within a few hours after application is recommended for maximum mycorrhizal colonization and microbial proliferation.

THIS PRODUCT MAY BE APPLIED AT A MAXIMUM RATE OF 4 GALLONS (38.24 LBS) PER 1,000 SQ FT PER YEAR.

TURF:

Use **SuperRootsTurf™** with 4 species of Endo Mycorrhizae.

Compatibility: This product is compatible in a tank mix with Subdue, Phosphite and Medallion fungicides for root rot treatments, and will not hinder the function of the biological components. Other combinations may be acceptable, but testing should be done to ensure the biological components are protected and other compatibility issues are addressed.

CAUTION: This fertilizer is to be used only on crops which respond to molybdenum. Crops high in molybdenum are toxic to grazing animals (ruminants).

WARRANTY: Plant Essentials, LLC warrants that this product conforms with the guaranteed analysis. No other warranty is expressed or implied including any consequential damage, injury, or loss due to the use of this product by itself or in combination with any other product. Any liability is limited to product replacement.



3944 South Bay Rd NE, Olympia, Washington 98516
www.plantessentials.biz • 360.491.0556

Years of extensive field use of this product have proven **SuperRoots™** to be a valuable, easy-to-apply product when used as directed in a treatment-based plant health care program. If plants do not respond, steps should be taken to determine if there are soil conditions that would hinder plant health. Some examples are: improper site location, improper soil pH level, poor planting and/or watering practices, the presence of root rotting disease pathogens or insects, and soil compaction.